

TECHNICAL REPORT DOCUMENTATION PAGE

1. Report No.: TX 99/7-4903-1	2. Government Accession No.:	3. Recipient's Catalog No.:	
4. Title and Subtitle: Right-of-Way Mowing Height Research: Literature Review		5. Report Date: February 1999	
		6. Performing Organization Code: TechMRT	
7. Author(s): John Borrelli, Cynthia McKenney, Ron E. Sosebee, and Richard Zartman		8. Performing Organization Report No.: 7-4903-1	
9. Performing Organization Name and Address: Texas Tech University Center for Multidisciplinary Research in Transportation Box 41023 Lubbock, Texas 79409-1023		10. Work Unit No. (TRAIS)	
		11. Contract or Grant No.: 7-4903	
12. Sponsoring Agency Name and Address: Texas Department of Transportation Research and Technology P. O. Box 5080 Austin, TX 78763-5080		13. Type of Report & Period Cover: Deliverable #1	
		14. Sponsoring Agency Code:	
15. Supplementary Notes:			
16. Abstract: The focus of this project is to develop practical information and recommendations on the relationship between height and frequency of mowing. There is concern that current mowing practices are not timed to maximize the viability of the mixture of plants found in highway right-of-ways. It is hoped that mowing frequency can be reduced in some locations to minimize right-of-way maintenance costs. This project is intended to determine the optimum mowing height with respect to maintaining a viable stand of acceptable vegetation, reseeding of annual plant materials especially native flowers and the frequency of required mowing. The optimum height must also be considered with respect to the total cost of maintenance of right-of-way vegetation. To accomplish objectives, plots will be established in four locations and responses measured to various treatments of mowing height and frequency of mowing. Measurements will be taken on plant population changes, morphological development, root depth, ground shading, and carbohydrate reserves in plants. In addition, data will be collected on the related issues of mowing height and mower speed versus rooster tails, and on the quantity and size of debris that might impact the choice of mowing height.			
17. Key Words: mowing height		18. Distribution Statement: No restrictions. This document is available to the public through the National Technical Information Service, Springfield, Virginia 22161	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 44	22. Price

This page replaces an intentionally blank page in the original.

-- CTR Library Digitization Team

RIGHT-OF-WAY MOWING HEIGHT RESEARCH:
LITERATURE REVIEW

by

John Borrelli, Cynthia McKenney
Ron E. Sosebee and Richard Zartman

Research Report Number 7-4903

Conducted for:
Texas Department of Transportation

by the

CENTER FOR MULTIDISCIPLINARY RESEARCH IN TRANSPORTATION
TEXAS TECH UNIVERSITY

February 1999

This page replaces an intentionally blank page in the original.

-- CTR Library Digitization Team

IMPLEMENTATION STATEMENT

The results of this project will yield the documentation necessary to verify or provide the scientific information needed to revise CMD's Roadside Vegetation Management Manual with regard to mowing height and frequency of mowing. A Research Report, Project Summary Report, educational materials, and an Internet document on the Texas Tech University's TechMRT Internet site will convey the findings of this project.

Prepared in cooperation with the Texas Department of Transportation and the U.S. Department of Transportation, Federal Highway Administration.

AUTHOR'S DISCLAIMER

The contents of this report reflect the views of the authors who are responsible for the facts and the accuracy of the data presented herein. The contents do not necessarily reflect the official view of policies of the Department of Transportation or the Federal Highway Administration. This report does not constitute a standard, specification, or regulation.

PATENT DISCLAIMER

There was no invention or discovery conceived or first actually reduced to practice in the course of or under this contract, including any art, method, process, machine, manufacture, design or composition of matter, or any new useful improvement thereof, or any variety of plant which is or may be patentable under the patent laws of the United States of America or any foreign country.

ENGINEERING DISCLAIMER

Not intended for construction, bidding, or permit purposes.

TRADE NAMES AND MANUFACTURERS' NAMES

The United States Government and the State of Texas do not endorse products or manufacturers. Trade or manufacturers' names appear herein solely because they are considered essential to the object of this report.

SI* (MODERN METRIC) CONVERSION FACTORS									
APPROXIMATE CONVERSIONS TO SI UNITS					APPROXIMATE CONVERSIONS FROM SI UNITS				
Symbol	When You Know	Multiply By	To Find	Symbol	Symbol	When You Know	Multiply By	To Find	Symbol
LENGTH					LENGTH				
in	inches	25.4	millimeters	mm	mm	millimeters	0.039	inches	in
ft	feet	0.305	meters	m	m	meters	3.28	feet	ft
yd	yards	0.914	meters	m	m	meters	1.09	yards	yd
mi	miles	1.61	kilometers	km	km	kilometers	0.621	miles	mi
AREA					AREA				
in ²	square inches	645.2	square millimeters	mm ²	mm ²	square millimeters	0.0016	square inches	in ²
ft ²	square feet	0.093	square meters	m ²	m ²	square meters	10.764	square feet	ft ²
yd ²	square yards	0.836	square meters	m ²	m ²	square meters	1.195	square yards	yd ²
ac	acres	0.405	hectares	ha	ha	hectares	2.47	acres	ac
mi ²	square miles	2.59	square kilometers	km ²	km ²	square kilometers	0.386	square miles	mi ²
VOLUME					VOLUME				
fl oz	fluid ounces	29.57	milliliters	mL	mL	milliliters	0.034	fluid ounces	fl oz
gal	gallons	3.785	liters	L	L	liters	0.264	gallons	gal
ft ³	cubic feet	0.028	cubic meters	m ³	m ³	cubic meters	35.71	cubic feet	ft ³
yd ³	cubic yards	0.765	cubic meters	m ³	m ³	cubic meters	1.307	cubic yards	yd ³
MASS					MASS				
oz	ounces	28.35	grams	g	g	grams	0.035	ounces	oz
lb	pounds	0.454	kilograms	kg	kg	kilograms	2.202	pounds	lb
T	short tons (2000 lb)	0.907	megagrams (or "metric ton")	Mg (or "t")	Mg (or "t")	megagrams (or "metric ton")	1.103	short tons (2000 lb)	T
TEMPERATURE (exact)					TEMPERATURE (exact)				
°F	Fahrenheit temperature	5(F-32)/9 or (F-32)/1.8	Celcius temperature	°C	°C	Celcius temperature	1.8C + 32	Fahrenheit temperature	°F
ILLUMINATION					ILLUMINATION				
fc	foot-candles	10.76	lux	lx	lx	lux	0.0929	foot-candles	fc
fl	foot-Lamberts	3.426	candela/m ²	cd/m ²	cd/m ²	candela/m ²	0.2919	foot-Lamberts	fl
FORCE and PRESSURE or STRESS					FORCE and PRESSURE or STRESS				
lbf	poundforce	4.45	newtons	N	N	newtons	0.225	poundforce	lbf
lbf/in ²	poundforce per square inch	6.89	kilopascals	kPa	kPa	kilopascals	0.145	poundforce per square inch	lbf/in ²

* SI is the symbol for the International System of Units. Appropriate

TABLE OF CONTENTS

Table of Contents

Technical Document Page.....	i
Title Page.....	iii
Implementation Statement.....	iv
Disclaimers.....	vii
Metric Table.....	viii
Table of Contents.....	ix
Introduction.....	1
Selection of Plots.....	2
Review: Defoliation of Grasses.....	4
Introduction.....	4
Literature Review.....	4
Growth Habit of Grasses and Affect or Defoliation on Plant Response.....	5
Carbohydrate Reserves.....	6
Defoliation Height and Frequency.....	7
Tiller Development.....	8
Management of Right-of-Way Vegetation.....	9
Introduction.....	9
Mowing Height.....	10
Plant Root Development.....	12
The Affect of Vegetation Height in Right-of-Ways on Snow Drifting.....	14
Introduction.....	14
Height and Porosity of Wind Barriers.....	14
Porosity of Vegetative Barriers.....	16
Simulation of Vegetative Wind Barriers Adjacent to Shoulder.....	17
Summary.....	18
Litter Review.....	19
Introduction.....	19
Review of Major Litter Surveys.....	19
Summary.....	22
References.....	23

Introduction

This progress report present the findings of several literature searches into the relationships between grasses and the height and frequency of mowing. In addition, it contains the description of the sites for the four sets of plots. The data bases searched included TRIS as well as traditional sources including the internet.

The first part of the report provides the location of the plots along with photo of the sites. The plots are located in Tahoka, Lufkin, Andrews, and Brady. Each site appears representative of a geographical areas of Texas and has representative vegetation for the area. Selection of the sites were made with the aid of TxDOT personnel.

The second part of this report contains the literature reviews concerning defoliation of grass by mowing, carbohydrate reserves in grasses, tiller development, mowing height, plant root development, litter, and vegetative wind barriers. Approximately 150 references are cited in the review. The information obtained from this review will help guide the research. As work progresses, additional citations will be found and information gathered on more specific topics.

Selection of Plots

Four sets of plots were selected at Andrews, Brady, Lufkin, and Tahoka. These plots were selected with the aid of Ms. Ann Finley for the Lubbock District, Mr. Kyle Moseley for the Brownwood District, Mr. Herbert Bickley for the Lufkin District, Mr. Robert Watts for the Odessa District. The purpose of the plots is to evaluate the response of grasses to different mowing heights and frequency of mowing. The purpose of the four locations is to allow for diversity of climate, precipitation, and vegetation to be included in the study.

The experimental design will be a split plot with four replications per location. The four locations were chosen near Andrews, Tahoka, Brady and Lufkin, Texas. The main treatment will be three mowing heights at 2, 4 and 8 inches. The secondary treatment will be frequency of mowing. Plots will be mowed once a year (October), twice a year (July and October), or three times a year (May, July, October).

Each plot will be 50 feet long, with a 20 foot border between individual plots. Plot width will be 8 feet to correspond with the swath of the rotary mower. Plots will be randomized with four replicates at each of the four locations. The plots will be located as near to the right-of-way fence as possible to minimize disturbance from unforeseen traffic. The plots will have signage of "no-mow zones" to prevent normal mowing operations from inference with the experimental mowing schedule.

Presented below are the descriptions of the plot locations including pictures of each site. There is no scheduled construction activity within the next 5years for any of the locations. The sites are representative of the conditions existing in each of the districts and should provide the vegetative and climate diversity needed for this study.



View looking south

The plots are in the right-of-way on US 385 and are located as follows: approximately 7.45 mi. north of the intersection of US 385 and SH 158 or approximately 3.17 miles north of the Ector and Andrews County lines. They are just south of the Windmill roadside park.

Figure 1.-Andrews County near Andrews.



View looking southwest



View looking northeast.

The plots are located on the south right-of-way of HY 190 approximately 3.9 miles northeast from the intersection of HY 377 and HY 190.

Figure 2.-McCulloch County near Brady.



View in center of plot.



View from southwest end.

The plots are approximately 1.5 miles from the junction of HY 2497 southwest on HY94. The plots are located on the north right-of-way. The gully is to be repaired.

Figure 3.-Angelina County near Lufkin.



View looking west.



View looking east.

The plots are approximately 6.5 miles west of Tahoka on HY 380. The plots are located on the south right-of-way.

Figure 4.-Lynn County near Tahoka.

Review: Defoliation of Grasses

Introduction

Plant response to defoliation has been emphasized because of the extensive use of rangelands as a forage source for both domestic livestock and wildlife. Knowledge of plant response to defoliation has been incorporated into various systems and prescriptions to effectively manage both vegetation and herbivores to minimize the detrimental consequences of grazing and to maintain plant and animal production on a sustainable basis (Briske and Richards, 1995).

Many studies on the effect of defoliation on plant vigor, productivity, and carbohydrate reserves have concluded that frequency, intensity, and season of defoliation could explain most of the plant response to defoliation (Dahl and Hyder, 1977). Consequently, the most favorable leaf replacement can be attained by scheduling the time (frequency) and degree (stubble height) of defoliation to stop the culm growth and promote new tillers (Hyder, 1972).

Grazing may be timed to primarily produce a strong tillering response, harvest a maximum quantity of nutrients per unit area, and improve forage quality for subsequent grazing. It may also be used to manipulate botanical composition, protect and improve adjacent range, or create some other desirable effects (Hyder, 1974).

Other desirable effects of planned defoliation could be practiced to afford protection against fire hazards along highway rights-of-way by maintaining a minimum stand of vegetation with the ability to regenerate after defoliation. Also, restoring an aesthetically pleasing environment is another desirable effect that could be considered.

Although there is a significant amount of literature on the effects of height and frequency of defoliating (mowing) native grasses, most of it does not relate defoliation to either the phenological stage or the physiological status of the plant. Therefore, this portion of the research project is designed to test the effect of height and frequency of defoliation and the time of defoliation on grass response to defoliation. A representative from tall, mid, and short grasses will be selected (to the extent possible) for evaluation.

Literature Review

In general, defoliation of range plants affect many related factors, and it seems unlikely that defoliation may effect one factor without simultaneously influencing the others. This literature review will address a few of the factors such as growth habit, carbohydrate reserves, and tiller development that are affected by frequency and intensity of defoliation throughout the growing season.

Growth Habit of Grasses and Affect of Defoliation on Plant Response

Most grasses exhibit morphological adaptations that permit them to endure partial defoliation. The apical meristem which is the source of new leaves is elevated by stem elongation; therefore, the removal of, or damage to, the apical meristem stops stem elongation and leaf expansion from that axis. Subsequent growth and leaf replacement is delayed until a new stem arises from the basal buds (Hyder, 1972). Removal of the apical meristem stimulates axillary shoot growth regardless of the stage of development of Arizona cottontop (*Digitaria californica* (Benth.) Chase) (Cable, 1982). The concept of apical dominance has been studied by several range scientists to explain the regulation of tillering in grasses, especially in response to defoliation. Removal of the apical meristem stimulates axillary bud growth and promotes tiller initiation. The stimulation of bud growth presumably occurs because the direct inhibitory influence exerted by the apical dominance is removed (Murphy and Briske, 1992).

Range plants which are most resistant to excessive grazing grow in a manner such that their apical meristems are inaccessible to grazing animals. These plants have the appearance of being culmless because of their extremely short basal internodes (often called short shoots) so that the elevated part of a leaf blade can be removed without stopping growth (Dahl, 1995). Defoliating perennial ryegrass (*Lolium perenne*) in the early vegetative stage (stem elongation or flowering stages) produced higher annual standing crop yields than treatments in which plants were defoliated during an intermediate stage of growth when the majority of stem apices had been elevated above the height of defoliation (Binnie et al., 1980). On the other hand, long shoots which elevate their apical meristem early in the growing season and, thus, are exposed to grazing animals always decrease in standing crop under continuous heavy grazing (Dahl, 1995). The grasses in which the growing points reached a height that permits removal of the apical meristem by grazing; e.g. switchgrass (*Panicum virgatum*) decreased as intensity of utilization (grazing) increased, but grasses with growing points at the ground level, e.g. blue grama (*Bouteloua gracilis*) and sideoats grama (*B. curtipendula*) usually increased their standing crop (Branson, 1953). Also, sideoats grama and little bluestem (*Schizachyrium scoparius*) exhibited the ability to tiller because they have many basal short buds along with adventitious roots that supply the needs of newly developed lateral branches, while Indiangrass (*Sorghastrum nutans*) and switchgrass, which have fewer basal buds, were less able to spread or even maintain growth under intensive grazing (Rechethin, 1956).

Continuous grass growth during the summer could be possible for both long and short shoot plants if proper management is practiced. Timothy (*Medicago sativa*) and brome (*Bromus sp.*) grasses both characterized by long shoots, tend to go dormant when growing points are grazed off too early. At later growth stages, the growing points must be grazed off to activate lateral buds in the root crown to initiate tiller development. While in orchardgrass (*Dactylis glomerata*) and big bluegrass (*Poa ampla Merr.sp.*) (grasses in which their growing points remain at ground level, or below), frequent grazing is necessary to protect the growing points, but a sufficient number of leaf blades must be left to keep the growing point supplied with adequate moisture to prevent drying

out (Heppner, 1961). However, clipping little bluestem, big bluestem (*Andropogon gerardi* Vitman), and Indiangrass at any time during the summer reduced yield, but clipping at 3.75-cm stubble height between floral initiation and anthesis was most damaging to the plant (Vogel and Bjustad, 1968).

Clipping little bluestem to 7.5-cm stubble height before internode elongation reduced production; but, clipping the plant to the same stubble height after internode elongation increased production. Herford (1951) also found that clipping to 5-cm stubble height in July increased production, but clipping to 10 and 15 cm on the same date decreased yield because the apical meristem was removed by the 5-cm clipping, but not by the 10- or 15-cm stubble heights (Jameson and Huss, 1958). Clipping the entire basal area of bluebunch wheatgrass (*Pseudoroegneria spicata* [Pursh] Scribn. & Smith) plants to a 7.6-cm stubble height during the mid-boot phenological stage and the inflorescence emergence stage caused 7.0 and 7.5% declines in live basal area, respectively (Clark et al., 1998).

Carbohydrate Reserves

The major premise for monitoring carbohydrate reserves is to provide an index of potential for regrowth based upon the assumption that these carbohydrates provide the major source of carbon for leaf regrowth. These labile carbon compounds are utilized for plant growth and maintenance when the photosynthetic capacity is limited as demonstrated by the reduction in carbohydrate reserves following defoliation (Coyne et al., 1995). In seasonal comparisons, an inverse relationship is frequently observed with carbohydrate pools increasing during periods of minimal growth and then decreasing during periods of rapid growth (Briske and Richards, 1994). Plants that replenish reserves rapidly after spring “draw down” and regrowth periods and minimize the portion of the growing season with low reserve status were least affected by defoliation and recovered rapidly from severe defoliation. These plants are characterized by a narrow V-shaped curve in TNC trends such as in blue grama (Menke and Trlica, 1981). Defoliation at any time of the year usually effects plant growth by affecting carbohydrate reserve level, however, plants were found to be more affected by defoliation at certain phenological stages than at others (Trlica, 1977). There was a direct relationship between average total carbohydrate levels in the autumn and the amount of regrowth after defoliation. The carbohydrate reserve level in the autumn appears to be a good indicator of defoliation effects during the preceding growing season for some semi-arid species such as needle-and-thread (*Stipa comata*) and Indian ricegrass (*Oryzopsis hymenoides*) (Trlica and Cook, 1971). Also, results indicated that defoliation of mountain plants such as needlegrass (*Stipa lettermanii*), beardless wheatgrass (*Agropyron inerme*), and rabbitbrush (*Chrysothamnus viscidiflorus*) early in the growing season when the carbohydrate reserves were low was more detrimental than defoliation late in the season when the carbohydrate reserves were high (Donart and Cook, 1970). Moreover, maximum plant vigor in relation to carbohydrate reserves depends upon reserve storage sometime at the end of the growing season for the semi-arid species Indian ricegrass and needle-and-thread (Coyne and Cook, 1970). Monthly defoliation decreased carbohydrate concentrations by about 50% and carbohydrate pools by 80%, respectively, within stems, leaves, and crowns of

spotted knapweed (*Centaurea maculosa* Lam.) (Lacey et al., 1994). Complete protection of perennial ryegrass (*Lolium perenne*) from grazing did not increase carbohydrate reserves accumulation when compared to the grazing treatments; however, non-grazing sometimes produced significantly lower carbohydrate concentrations than grazing treatments (El Hassan and Krueger, 1980). Defoliation to short stubble heights led to greater reductions in non-structural carbohydrates. While, with taller stubble heights, stand composition, and non-structural carbohydrate levels improved in big bluestem (Forwood and Magai, 1992).

Both defoliation and drought stresses were found to effect carbohydrate reserve levels in two cool season grasses; crested wheatgrass (*Agropyron desertorum* Fisch. ex Link) Schult.) and bluebunch wheatgrass. Plants exposed to prolonged periods of drought plus defoliation once a year at 5- to 7-cm stubble height may have rapid initial regrowth upon alleviation of both stresses because high amounts of total nonstructural carbohydrates may have accumulated in their storage organs during stress (Busso et al., 1989).

Defoliation Height and Frequency

There is a relatively large amount of information in the literature concerning the effect of height and frequency of defoliation, but almost none of the literature relates the response of defoliation to the phenological stage of the plant. Therefore, only a few references address the relation of defoliation height and frequency to phenological stages.

The effect of defoliation height and frequency on tiller initiation is difficult to generalize; it is easily confounded with phenological development and the seasonal progression of environmental variables (Briske and Richards, 1995). The timing of defoliation in the spring appeared to be the most important factor governing basal crown bud production and regrowth for phalaris (*Phalaris aquatica* cultivar Sirolan) in the following summer (Hill and Watson, 1989). Also, net growth rate declined significantly as the time of clipping (at 10 cm) was delayed July 1 through the April-October growing season in the tallgrass prairie (Gillen and McNew, 1987). It is recommended that pinegrass (*Calamagrostis rubescens*) be grazed for a short time while it is actively growing (early in June) and then later when quiescence is well established (August) to maintain its vigor (Darryl et al., 1980). Moreover, rough fescue (*Festuca scabrella* var *campestris* Rydb) is highly sensitive to grazing when the plant is growing. It has been suggested that the greatest benefit from rough fescue grasslands may be derived by grazing in the fall or winter (Willms, 1991). Total annual forage production on plots clipped during the boot stage was generally lower than on plots clipped during the vegetative or late-flowering stages of development in crested wheatgrass (Miller et al., 1990).

Forage harvested during the growing season was greatest from plots harvested three or more times in the fall and least when big sacaton (*Sporobolus wrightii*) was harvested in the spring, spring/early summer, and spring/midsummer (Haferkamp, 1982). While defoliating little bluestem at 7 cm, a single defoliation in June or July during the

growing season produced the optimum dry matter yield, tiller weight and number, and bud number when compared to a single August defoliation, or two or three early or late defoliation treatments (Mullahey et al., 1990).

Defoliation early at the three-leaf stage of white clover (*Trifolium repens*) decreased the number of growing points by 32% and stolon weights and lengths by 50% compared with delaying defoliation until the nine-leaf stage (Hayes and Williams, 1995). Cutting creeping bluestem (*Schizachyrium stoloniferum* Nash.) at 10 cm above the soil surface significantly reduced dry matter, but there was no significant drop in yield when plants were cut at a 20-cm stubble height (Kalmbacher et al., 1981). Defoliating oats (*Avena strigosa* cv. saia) and ryegrass (*Lolium perenne* cv. Kangaroo Valley) every 6 weeks at a 5-cm stubble height produced the greatest yield (Lowe and Bowdler, 1988). Clipping tall wheatgrass (*Agropyron elongatum* (Host) Beau. 'Jose') every 4 weeks produced greater herbage than clipping every 1 or 2 weeks (Undersander and Naylor, 1987). The optimum time and frequency of defoliation for annual dry matter, and bud and tiller number was a single June or July defoliation for sand bluestem (*Andropogon gerardi* var. *paucipilus*) and a June-August defoliation for prairie sandreed (*Calamovilfa longifolia* (Hook.) Scribn.) (Mullahey et al., 1991). Interrupting growth of northern wheatgrass (*Agropyron dasystachyum*) by defoliation any time during the growing season disrupts production potential that year and the following year (Kowalenko and Romo, 1998). Average yields of northern wheatgrass were higher when the first defoliation was delayed until August and repeated at 6-week intervals (Zhang and Romo, 1994). Dry matter production was greatest under the 4-week/7-cm defoliation regime and least under the most severe defoliation regime (2-week/2-cm) in phalaris and orchard grass (Hill and Watson, 1989). Prairie grass (*Bromus wildehowii* Kunth cv. Matua) tiller numbers were increased by less frequent defoliation and were greater under the 8-cm than the 3-cm defoliation height (Bell and Ritchie, 1989).

Tiller Development

The ability to tiller or branch from axillary buds without the apical meristem being removed is considered an indication of an efficient forage producer. Richards et al. (1988) found that rapid production of a new photosynthetic canopy immediately following defoliation is critical for response of species which are tolerant of herbivory, especially when defoliated plants must compete with non-defoliated neighbors. Crested wheatgrass is an outstanding example of a species that tillers profusely following defoliation (Dahl, 1995). Rates of tiller growth on defoliated plants may occasionally exceed those of non-defoliated plants. In crested wheatgrass, grazed tillers had higher relative growth rates when grazing occurred prior to internode elongation in the spring than tillers of non-grazed plants (Briske and Richards, 1995). Also, Burzlaff et al. (no date) found that tiller development in blue grama depended on the reproductive culm of a shoot reaching maximum elongation.

Management of Right-of-Way Vegetation

Introduction

Maximizing the effectiveness of controlling vegetation on the highway right-of-way is of constant concern throughout the State of Texas. Existing vegetation on the roadsides needs to be maintained in a healthy status for safety and aesthetic considerations as well as to prevent soil erosion with the least amount of inputs. Many factors and interactions influence the vegetation along the highways in Texas. Four diverse geographic regions of Texas have been selected to be evaluated in this study. These four areas were selected to develop practical information on the relationship between height and frequency of mowing and the viability of maintaining vegetation within highway right-of-ways. One of our first priorities was to develop an understanding of the state-of-the art with respect to mowing height effects on plant density, rooting depth, and stand viability. This literature review will concentrate on the management problems of highway vegetation. The literature was found in a diversity of sources that sometimes directly address the problems in question regarding roadside vegetation practice. Many times these data must be interpreted from other literature with similar studies using different methods or species.

In The Highway and the Landscape, Snow presented the general topics of the interrelationship of highways and landscapes (Snow, 1959). The book has specific chapters devoted to the concepts and challenges of having roads in the landscape. Specific chapters address the complete highway (Bugge and Snow, 1959), fitting the highway into the landscape (Cron, 1959), preserving the scenic qualities (Johnson, 1959), the politics of road building (Ingraham, 1959). The most relevant chapter to this particular project is entitled "The functional uses of plants on the complete highway" (White, 1959). In that chapter, White discusses the utility, safety, and public acceptance of plants along the highways. The concerns written about in 1959, are still relevant to highway right-of-way maintenance today.

There are numerous general turfgrass management references available in book, journal article, nonrefereed article and web site formats on the subject of management of roadside vegetation. General text references include Beard, 1973 (Turfgrass Science and Culture); Duble, 1996 (Turfgrasses: Their Management and Uses in the Southern Zone); Madison, 1971 (Practical Turfgrass Management); Schroeder, 1996 (Turf Management Handbook); and Turgeon, 1999 (Turfgrass Management). Cogburn (1995) TxDOT mowing guidelines would be an example of a refereed journal article on the subject. Nonrefereed articles or handbook type publications would be those such as Turfgrass Maintenance and Establishment: A Student Handbook (Penn. State Univ., 1968); Desirables and Weeds for roadside Management--a Northern Rocky Mountain catalogue (Meier and Weaver; 1977); Giving Nature a Hand booklet (TxDOT, 1999); Roadside Vegetation Management Manual: A volume of the Infrastructure Maintenance Manual: Manual Notice 93-1 (TxDOT, 1993); and Roadside Vegetation Management Manual: Manual Notice 96-1 (TxDOT, 1996). World Wide Web formats are now rapidly supplanting the traditional manuals as methods to disseminate the roadside management

information. Several of the web addresses are as follows:
<http://www.dot.ca.gov/hq/LandArch/> and
<http://plants.usda.gov/plantproj/plants/index.html>.

Mowing Height

Plant growth is a function of many environmental and management components. The environment is a function of the climate of the area and the soil resources of that area. Climate cannot be controlled, in general, other than by irrigation and/or drainage that are not pertinent to this document. The climatic factors are evaluated in this project by the selection of four different geographical regions to conduct this study. Soil resources are generally a function of the native resources and the history of the highway's construction at that location. These resources can be amended by the addition of fertilizer or liming materials. As these practices are generally not considered to be justifiable due to the acreage encompassed, these factors will not be included in this review. The factors of soil compaction, mowing height and mowing frequency on vegetation growth and the detrimental influences of roadway management practices will be discussed.

The primary management tool available to the Texas Department of Transportation (TxDOT) is mowing. Parameters such as mowing height, mowing frequency and mowing time can be controlled and prescribed. Many publications have been devoted to explaining these phenomena and there is not one solution to all situations. A review monograph (Turfgrass, Waddington et al., 1992) reviews the general physiology; soils and water; and management of turfgrass. The monograph has physiology chapters related to ecological aspects (Watschke and Schmidt, 1992); energy relations and carbohydrate partitioning (Hull, 1992); salinity (Harivandi et al., 1992); and physiological effects of temperature stress (DiPaola, 1992). The soils and water chapters relate to nutritional requirements (Turner and Hummel, 1992) and water requirements (Kneebone et al., 1992). The management chapters address energy conservation (Busey and Parker, 1992); integrated pest management (Bruneau et al., 1992); and plant growth regulators (Watschke et al., 1992). This review monograph does not, however, address the management implications of mowing to any great extent.

Mowing is the most fundamental practice utilized to maintain the aesthetics and safety functionality of the highway right-of-way. While herbicides may be incorporated into the overall management program (McCully and Bowmer, 1971), mowing provides a uniform surface for viewing traffic and any potential hazards to driving. Since mowing defoliates the plants, many physiological changes of the vegetation occur. While maintaining visibility, mowing must not deplete the carbohydrate reserves of the plant material to such an extent that the vegetation dies. The criteria for selecting the mowing height and frequency are physiological condition and growth habit of the vegetation as well as safety considerations such as visibility and flammability.

The choice of mowing height is dependent on the physiology and the timeliness of the mowing (Clapp, et al., 1964; Cogburn, 1995; Davis, 1967, Kneebone, 1966).

Mowing has been reported to increase succulence of the above ground portions of the plant and increase the quantity of chlorophyll per unit area (Madison, 1962). Other reports indicate decreased leaf width with mowing (Harrison, 1931; Mitchell, 1955; Mitchell and Coles, 1955; and Robertson, 1933). Closely mowing turfgrass generally produces a finer, denser foliage, but has less underground roots (Biswell and Weaver, 1933; Turgeon, 1999).

Mowing vegetation short has been reported to make the vegetation more susceptible to heat and water stress (Salaiz et al., 1995). This susceptibility is probably due to decreased carbohydrate synthesis and storage (Bukey and Weaver, 1939; Dodd and Hopkins, 1958; Everson, 1966; and McCarty and Price, 1942). Practices such as strip, modified full and full width mowing are all management tools to be utilized in highway right-of-way management. Heat stress induced problems would occur more often with the late spring and summer mowing than in the final clean cut that occurs after the first frost in the fall. Razmjoo et al. (1996) report that grasses improve when mowing height is increased. Fluck and Busey (1988) state that low mowing heights requires greater amounts of fertilizer input to give the same benefit as allowing the mowing height to be raised.

Mowing height dramatically affects the root development (Biswell and Weaver, 1933; Burton et al., 1963; Crider, 1955; Everson, 1966; Goss and Law, 1967; Graber and Ream, 1931; Harrison, 1931; Harrison, 1934; Harrison and Hodgson, 1939; Juska and Hanson, 1961; Law, 1966; Madison, 1962; Robertson, 1933; and Sprague, 1933). Carter and Law (1948) and Branson (1956) observed that defoliation had a negative effect on root development. Branson observed that herbage removal of range grasses decreased yields of tops and roots as removal intensity increased. Defoliation more dramatically influenced root development than shoot development. The effects of root production were observed by Branson (1956) on blue-bunch wheatgrass. Root production of the control plants was observed to be more than 100 times greater than the clipped plants. Carter and Law (1948) observed similar results when they clipped other perennial grasses at different intervals. They reported that clipping at 30-day intervals severely retarded root and top growth of all species. Graber and Ream (1931); Harrison (1931); Harrison (1934); Harrison and Hodgson, (1939); and Hughes, (1937) all reported decreased rhizome growth as a function of mowing warm season grasses.

Roadside vegetation is not only important for safety and aesthetics, but also in the control of erosion. There are many general references that cover the topic of vegetation and soil erosion (Crovetto, 1996; Hudson, 1995; Larson, et al., 1983; Schwab et al., 1996; and Troeh et al., 1980). For Texas, there are several relevant articles that evaluate erosion control on highway right-of ways (McCully and Bowmer 1967,1969). There are also environmental and economic considerations of maintaining an aesthetically pleasing highway right-of-way (Pimental et al., 1995). Vegetation absorbs the kinetic energy of raindrop impact and buffers the soil from this erosive action (Krenisky et al., 1998; Lattanzi et al., 1979). Velocity of runoff is also diminished with good surface cover (Hart, 1974). Vegetation also acts to minimize the wind impact on the soil surface in those areas susceptible to wind erosion (Skidmore and Siddoway, 1978).

Plant Root Development

The first environmental factor to be discussed will be the influence of soil physical factors on plant root development. The physical properties of soil bulk density, compaction, texture, shear strength and structure influence the rooting habits of plants. High bulk density soils (bulk density is soil mass divided by total dry soil volume) dramatically limits the rooting of plants growing along highways. Phillips and Kirkham (1962) studies indicated that densities less than 1.3 g/cm^3 were of little restriction to plant root growth. As densities increase, the root length density (length of root per volume of soil) decreases linearly. Soil compaction, which directly decreases pore volume has been shown to decrease root distribution. Phillips and Kirkham (1962) and Schuurman (1965) report that soil compaction reduced root growth by reducing the pore volume and increasing the mechanical impedance. Meredith and Patrick (1961) report the use of farm equipment such as the tractors that mow the roadsides can cause compaction to such an extent that roots cannot penetrate the soil. Schuurman (1965) determined that whether a root was capable of penetrating a compacted soil depended both on the root's ability to penetrate the pore and the force necessary to push the soil particles aside. The mechanical obstruction offered by the cohesion of the soil particles was one of the main components to root development. Roots must push the soil particles aside to widen a pore enough to permit penetration.

Soil texture plays a major role in the penetration by the root. In moderately fine textured soils, the root development was decreased 40% between the least and most compacted zones (Meredith and Patrick, 1961). They reported that for medium textured soils, root penetration ceased when large pores were less than 2% of the soil matrix. Meredith and Patrick (1961) reported that even though large pores may be destroyed by compaction, if aeration is adequate, the soil is generally plastic enough to permit root penetration. Their field observations indicated that the effects of compaction was usually less permanent in a fine textured soil than in medium textured soils. In Texas, Gerard et al. (1972) reported that root elongation in clay soils decreased with increasing soil strength, particularly for soil strength's exceeding 1 MPa. For moderate strengthen soils, they reported better root penetration than for either low or high values of mechanical impedance soils. High root length density was promoted along the deformed zones or cracks in the dry soil. Taylor and Burnett (1964) observed that soil strength alone controlled the growth of roots through moist soil. They observed that soil strength of 2.5 to 3 MPa at field capacity prevented root penetration but roots will grow through a layer of soil at 1.9 MPa at field capacity. Barley (1963) reported those soils resist local deformation caused by roots when the soil strength is too great.

Hendrickson and Veihmeyer (1931) and Portas and Taylor (1976) determined distribution of roots as influences by soil-water content. Hendrickson and Veihmeyer (1931) reported that roots would not grow in soils drier than 1.5 MPa. Portas and Taylor (1976) observed that as soil water potential decreased from -0.6 to -90 MPa, root length decreased. They further reported that roots responded rapidly when the water content was increased.

Type of soil structure also influences the distribution of roots. White and Lewis (1969) evaluated the effects of soil structure on the distribution of several native grasses. They attempted to determine the effects of structure in dense, clay textured soils. They were able to correlate root system growth habits with soil structure by knowing the type of root system associated with the various grass species. The dense clay soil had larger more compact structural peds when dry than when wetter. Nearly dry ped aggregate size determined the width of the desiccation cracks in the soil. With small peds, the cracks were more numerous and had narrower widths than if the peds were larger. The size of the finest ped aggregate was an important factor as far as the root growth was concerned. Larger peds bordered by irregular cracks in the dense clay soils caused more root damage than the smaller peds of the clayey soils.

White and Lewis (1969) further observed that the shape of the soil structural unit was a major influence on root distribution. They determined that roots were stretched most frequently along prismatic structures where vertically oriented cracks form. Horizontal cracks, which were usually narrow due to the overlying weight, tended to counteract prism formation. Vertically growing roots cross fewer cracks than roots growing horizontally. Plants that have a vertical rooting habit, should be able to maintain a more stable population than those plants with a horizontal rooting habit. This difference would be due to their ability to adapt to soil structures with a vertical orientation. White and Lewis concluded that soil structures determined the distribution of grass species in clay soil by permitting better utilization of water.

A factor of equal importance to highway right-of way vegetation is the soil chemical properties and amelioration of these problems or corrections of deficiencies. Again, there are many general references that address these problems but, for the most part, are not typically utilized in right-of way vegetation management (Follett et al., 1981; Tisdale and Nelson, 1975; and Troeh and Thompson, 1993). Only in the most drastic cases in which a severe soil problem is encountered, are remedial actions taken (Thompson and Zartman, 1997).

There are also interactions between the fertility of the soil and mowing height. Harrison (1931) reported that different clipping heights affected the amounts of roots produced. In those grasses most susceptible to damage by mowing, it was not the cutting off the growth buds that was detrimental to growth. It was the gradual diminished carbohydrate reserves that lead to starvation beyond which the plants could not maintain themselves. Nitrogen fertilizer was reported to alleviate this problem to some limited extent (Razmjoo et al., 1996). Mortimer and Ahlgren (1936) not only evaluated fertility and mowing height, but also examined the irrigation effect. Irrigation may be used to modify the detrimental effects of mowing, but is generally not used along roadsides.

The Affect of Vegetation Height in Right-of-Ways on Snow Drifting

Introduction

The primary purpose of vegetation on the side-slopes of highways and in the right-of-ways is to control soil erosion and for the stability of the side-slopes (Hunt and Deschamps, 1995). Vegetation, however, can act as a wind barrier and can create drifting on the lee side of the barrier. Vegetation is commonly used as a wind barrier (Black et al., 1971). When vegetation acts as a wind barrier in the right-of-way of a highway, it can create a snow drift that extends on to the driving lanes creating a hazardous condition. The height of the vegetation after the final mowing is the height of the vegetative barrier which will create a snow drift during a wind-snow event.

The mowing height is related to the type of vegetation (primarily grass) desired to be maintained in the right-of-way. If the vegetation is cut relatively high (7 to 9 inches), the medium and tall grasses have an opportunity to establish and be part of the plant community in the right-of-way. If the vegetation is mowed short (~2 inches), short grasses will predominated and the medium and tall grasses will tend to disappear from the plant community. Thus, the decision of the Vegetation Manager on the height of mowing will dictate both the type of vegetation that will predominate in the right-of-way and the height of the vegetative wind barrier which will exist during the winter snow season.

Provided below is the relationship between barrier height and porosity with respect to snow drifts on the lee side of the wind barrier. The understanding of wind barriers and drifting have progressed allowing for reasonable predictions of snow drifts caused by both vegetative and man-made wind barriers.

Height and Porosity of Wind Barriers

Wind barriers are important for controlling both the movement of snow and the movement of soil (Greb, 1980; Skidmore and Hagen, 1977). Vegetative wind barriers have been recognized as being effective for reducing soil erosion and trapping water in the drifts that develop on the lee side of the barriers (Black et al., 1971). The main factors affecting the size and shape of the lee side drifts are the height and porosity of the barrier.

Borrelli et al. (1989) developed a mathematical relationship between the height, length of protection, and porosity for wind barriers. While the Borrelli et al. (1989) were concerned with soil erosion, the relationships can be applied the development of snow drifts behind a wind barrier. According to Greeley and Iversen (1985) the behavior of wind-blown dry snow is the same as sand. Greeley and Iversen (1985) stated that the average diameter of drifting dry-snow is 0.15 to 0.20 mm or with in the particle size range of dune sand.

A wind barrier causes a loss of momentum in the air stream. Because air at atmospheric pressure is essentially an incompressible fluid, a loss of momentum implies a reduction in wind speed. The reduction of wind speed causes the snow to deposit and form a drift. The greatest reduction in wind speed occurs approximately 6 barrier heights (6H) downwind from the barrier (Tabler, 1980). The wind profile is essentially reestablished to ambient conditions at approximately 26 times the height of the barrier downwind from the barrier.

Tabler (1980) developed an equation that predicted the length and height of drift for saturation conditions—the largest drift possible for the given porosity and height of the wind barrier. The equation developed by Tabler (1980) is

$$\frac{y}{H} = 0.13 + 0.402 \frac{x}{H} - 0.0602 \left(\frac{x}{H} \right)^2 + 0.003691 \left(\frac{x}{H} \right)^3 - 1.0854 \times 10^{-4} \left(\frac{x}{H} \right)^4 + 1.2498 \times 10^{-6} \left(\frac{x}{H} \right)^5 \quad \frac{x}{H} < 26.6 \quad (1)$$

where H = height of wind barrier (ft)
y = depth of snow (ft)
x = length of snow lee drift (ft)

Borrelli et al. (1989) developed a formula that predicted the leeward wind velocity downwind from the wind barrier. The formula developed by Borrelli et al. (1989) is

$$\frac{U_L}{U_o} = 1 - A e^{-0.0876 \frac{x}{H}} \quad (2)$$

where U_L = the leeward wind velocity at a height of 0.5H for location L (ft/sec)
 U_o = the upstream wind velocity at a height of 0.5H (ft/sec)
H = height of the barrier (ft)
x = the distance downstream from the barrier (ft).

The coefficient A is a function of porosity of the wind barrier. The equation for A is

$$A = 1.217 - 4.81 \times 10^{-3} P - 7.39 \times 10^{-5} P^2 \quad (3)$$

where A = the coefficient A in equation (2)
P = porosity of barrier in percent.

The formula developed by Tabler (1980), equation (1), gives essentially the same length of drift as the formula developed by Borrelli et al. (1989), equation (2), but in slightly different form. The two equations produce the same length of drift for conditions where the wind downwind of the barrier is 90 to 95 percent of the ambient wind speed or

the wind speed on the windward side of the wind barrier. As the wind speed approaches ambient wind speed (approximately 90 to 95 percent of ambient), the wind can now transport the snow and deposition ceases.

Equation (1) is for a wind barrier with 50 percent porosity, for a vertical-slat fence, and for the drift at saturation conditions (maximum drift size). This information can be used to determine the wind speed reduction existing at the terminal end of the snow drift. Using this information the ratio of U_L/U_o is 0.92 for x/H of 26.5. Tabler (1980) states that for the wind speeds encountered in nature, the drift shapes are independent of the wind speed. This is fortuitous because wind speed does not need to be factored into the equations to predict the length and depth for the maximum drift that can develop downwind from a wind barrier.

If we assume that the drift will not develop for a ratio of U_L/U_o greater than 0.92, then equation (2) can be used to estimate the length of a drift for wind barriers of different heights and different porosities. This assumption can be justified because sand and snow being transported by wind are essentially the same size and develop the same drift patterns (Greeley and Iversen, 1985; Pugh and Price, 1954). Furthermore, the length of the lee side drift is independent of wind speed for wind speeds of concern (Tabler, 1980; Woodruff and Zingg, 1952).

Porosity of Vegetative Barriers


Raine and Stevenson (1977) used vertical slats to make wind barriers of different barriers. Raine and Stevenson (1977) tested the wind barriers in the field and in a wind tunnel. The wind approached the barriers at right angles. Other researchers have used vegetative strips and have measured the porosity of the barriers. Hagen and Skidmore (1971) determined the average porosity for pampagrass to be 73 percent, a single row of sudangrass 49 percent, a double row of sudangrass 34 percent, a single row of grain sorghum 60 percent, a double row of grain sorghum 55 percent, and forage sorghum 58 percent. These plants are relatively tall and are large stemmed plants. Based on drag coefficients, Hagen and Skidmore (1971) stated that narrow plant windbreaks have relatively uniform porosity and drag coefficients similar to vertical-slat fences. One could conclude that a row of Johnsongrass skipped by a mower would be similar to forage sorghum. Furthermore, Moysey and McPherson (1966) stated that the shape of a barrier has less effect on leeward velocities than barrier porosity. While Johnsongrass may not have the same profile as sorghum, it should produce similar drifts providing the porosities are the same.

Black and Siddoway (1971) used tall wheatgrass barriers to trap snow in agricultural fields. The wheatgrass was planted in double rows with an estimated porosity of 40 percent. The effective height, H , was approximately 2.5 feet. The snow drift (if assumed to be maximum or at saturation) was $25H$ giving a U_L/U_o of 0.9 according to equation (2). Wheatgrass has relatively small stems and leaves in comparison to sorghum or sudangrass and is similar in structure to most grasses predominating in the right-of-ways of Texas highways. A double row of grass is

probably very similar to the grass immediately adjacent to the shoulders of a highway. Thus, the wind barrier created by grass immediately adjacent to a shoulder would not have a porosity any greater than 40 percent for grass cut uniformly in height.

Simulation of Vegetative Wind Barriers Adjacent to Shoulder

Presented in Figure 1 are the length of drifts for various height of barriers and barrier porosity. The height of barrier is the most important factor in determining the length of drift for porosities between 0 and 50 percent. For porosities greater than 50 percent, both the height of barrier and the barrier porosity are important factors.



For a good stand of vegetation adjacent to the shoulder of a highway, the porosity can be assumed to be 0 percent. This would be especially true for short vegetation (< 5 inches). For taller vegetation, there is significant porosity in the upper portion of the canopy and the overall porosity is probably near the values reported by Hagen and Skidmore (1971) for double rows of sudan grass or between 20 and 40 percent.

If rooster tails exist adjacent to the shoulder, the porosities would be similar to those reported by Hagen and Skidmore (1971) for single and double rows of sorghum and pampagrass. For a rooster tail that is 12 inches in height and with an assumed porosity of

75 percent, a drift of approximately 18 feet could result for a significant snow-wind event. This would potentially create a drift that could encroach onto the driving lanes.

Given equations (1) and (2) and a reasonable estimate of the porosity of the vegetation, one can estimate the length of the potential drift. The length of the drift provides an assessment of the hazard that could result from selecting the mowing height adjacent to the shoulder of the road. It also provides a measure of the risk created by rooster tails adjacent to the roads. This assumes that the highways are designed such that the wind clears the snow from the driving surface as recommended by Pugh and Price (1954).

The mowing height of the grass in the right-of-way more than 30 feet from the driving lanes will not add to the risk of drifting. Any drifts created by the vegetation would not reach into the driving lanes. One should be cognizant that if the vegetation is 7 inches in height, it would require a snow fall of greater than 7 inches or an accumulation of more than 7 inches before there would be any drifting potential.

Summary

Between the literature for wind breaks for wind erosion of soil and for wind breaks to control the drifting of snow, there is substantial knowledge concerning the effectiveness of vegetative wind breaks to control drifting of snow (Borrelli et al., 1989; Hagan and Skidmore, 1971; and Tabler, 1980). The most important parameter needed for vegetative wind breaks is the porosity of the vegetation. Black et al. (1971), Hagen and Skidmore (1971), Moysy and McPherson (1966), and Raine and Stevenson (1977) all reported porosities for vegetation including single-row and double-row strips of vegetation. The equation developed by Tabler (1980) and Borrelli et al. (1989) allow the prediction of the length of the potential drift and the depth of snow if needed. The body of knowledge appears scientifically sound and can be used for decision making without any great risk.

Litter Review

Introduction

Overall, the search of literature for documented sizes of litter in the right-of-way for Texas proved fruitless. In the literature found, it did not appear to be any significant quantities of large litter objects to pose harm to either the operator, the mowing unit, or a passerby (Syrek, 1986). Previous studies generally documented a large quantity of aluminum cans, small paper objects, and plastic wrappers with little or no count of larger, less frequent objects (auto parts, construction materials, etc) (Finkner, 1969; Syrek, 1986; Syrek, 1989; Syrek, 1990; and Vesilind, 1976). In no studies were there found to be significant amount by composition of large bulky items greater than the data from the survey of litter in California which reported approximately 38% of all litter on the roadside consisted of possibly harmful or detrimental objects (Syrek, 1989). It can be speculated that through the combination of strong litter awareness programs such as Texas Adopt a highway Programs, litter amounts in the state of Texas are lower than the majority of other states (Syrek, 1990).

Review of Major Litter Surveys

Several major litter surveys and Studies were identified during the literature background search. Of these studies, the 1969 National Study of the Composition of Roadside Litter contributed largely to the viewpoint of litter composition of right of ways (Finkner, 1969). Within this study, the Highway Research Board instigated a project to "obtain objective and adequate information regarding the composition and nature of litter scattered along the primary rural highways in the United States." Underlying this study was the main goal to gather and utilize data concerning the composition and nature of litter in the United States. Low participation throughout the nation led to the involvement of only 29 states, with Texas being one of the participants. Of the information gathered from these states, several generalized observations of litter composition were made.

A composition stratum of litter outlined that approximately 13% of the litter consisted of such items of concern as tires, lumber and other unclassified large objects, while 16% of the items were cans, and 6% was composed of bottles of various types and 59% was represented by the presence of paper items. Of this study, other generalizations such as the conception that there was a general trend of increasing litter volume with an increasing number of lanes. State by State reports were given, and of our concern, Texas' composition data along the right of ways were reported as follows composition:

Paper Items	48.59%
Cans	27.12
Number of Misc.	15.81
Bottles and Jars	5.33

The miscellaneous items included such items of interest as lumber or construction items, auto parts and accessories, tires and other unclassified objects, all of which would present a serious concern to mower safety if a higher percentage per total litter items. Information provided by Mr. Robert Watts indicates that the current TxDOT policy is to collect all large objects as encountered by TxDOT maintenance personnel. This practice essentially reduces the potential of encountering large objects in the process of mowing to a very low probability.

Another contributing data source to the literature search was Syrek (1986). Since the data are most current, and the survey originated from Texas, these data can be easily correlated to current Litter conditions along the right-of-ways. Although a major emphasis was placed on the actual change and reduction in litter volume rather than individual litter object compositions, this information still represented positive and valuable information. An overall reduction of 29% in litter was determined to have been achieved or "the largest of any 13 year-old trend comparisons surveys performed by the Institute for Applied Research." The litter counts described were divided into two part with the first being to compare the data with prior surveys of litter (considered to be a calibration process). The second 'count' governed the tabulation of four divisions of litter: cans, bottles, other deliberate litter (convenience products and packaging) and accidental litter. The considerable reduction in annual litter was determined to have been contributed to by the strong and intensive implementation of anti-litter programs such as public advertising and adopt a highway programs. Other highlights of the Texas survey suggested that of the actual litter along the roadsides, 34% of it could be seen by a pedestrian walking along the shoulder at a brisk pace. This study also examined other areas of interest such as the likeliness of one age group to litter more than another and what type of vehicle was the largest litter contributor, most of which were irrelevant to the concern of the current research project on the height to mow grass in the right-of-ways.

There was a similar survey of litter in California (Syrek, 1989). Syrek (1989) mainly concentrated on the change in beverage containers with in a years time span, and also touched on the composition breakdown of roadside litter. The Department of Conservation findings stated that by compositions (%), take-out food containers, cups, napkins, etc. composed the largest single product group found in the study. A decrease of 72% of bottles and cans from the previous year was noted following the introduction of the Recycling Act. The general breakdown of the estimated material compositions of California litter is listed below by percent composition:

Possible Significant items (percent)	
Wood lumber, pallets, signs	19.35
Rubber tires, Molding, Seals	14.10
Ferrous metals	2.6
Nylon rope, plastic pipe, other	1.02
Total	37.07

<u>Non-hazardous items (percent)</u>	
Glass bottles	23.20
Paper	16.26
Other	12.12
Aluminum cans	6.11
<u>Plastics</u>	<u>5.23</u>
Total	62.99%

In contrast to the surveys discussed earlier in other states, Syrek (1989) identified possible harmful objects comprise slightly under 40% of all litter items.

Another important and significant state study originated from the Louisiana Litter Survey of 1990 (Syrek, 1990). This study covered the collection of data from 110 locations around the state of Louisiana, documenting such instances as indiscriminate dumps. According to Syrek (1990), the largest single product identified in the group of litter in Louisiana was found to be take-out packaging, cups, napkins, plates, bags, etc., composing a total of 62% by composition. General data were collected which stated the occurrences of vehicle parts, and debris, construction materials, and wood pieces, comprising an approximate 10 percent (by composition) of all litter identified during the survey. An approximately 1 percent decline in food packaging in Louisiana and other states was noticed, while a 1-2 percent increase in construction material/wood materials was noticed for 1986. Embedded within this study was a comparison of Louisiana's "Indiscriminate Dump Encounter Rates" with those of other states. This information could then be used to draw a conclusion to the frequency per state to have an occurrence of a so called indiscriminate dump, or litter objects which would be of concern when mowing on the right of ways.

The data mentioned is listed below in terms of number of indiscriminate dumpings per 1000 miles of driving:

Louisiana (1990)	30.4
Hawaii (1988)	28.2
Florida (1989)	23.8
Pennsylvania(1984)	15.5
New York(1984)	13.9
Oregon (1986)	7.6
*Texas (1989)	5.6
California (1985)	3.9
Washington (1987)	1.7
Nebraska (1985)	1.0

It should be noted that Texas has a considerably low value of indiscriminate dumps per 1000 miles. It should also be noted that Syrek (1990) visited Texas and stated in the report that there was a low level of indiscriminate dumps in Texas. The contributing factors were speculated to be a strong public broad cast of the "Don't Mess with Texas" propaganda either by sign, TV, or radio.

One final relevant state survey was examined—a study conducted in Michigan (O'Toole, 1979). The litter study counted items larger than a cigarette package, in a effort to exclude smaller items such as gum wrappers or pull tabs. The method for classification centered around the division of 5 items: paper, plastic, cans, bottles and miscellaneous objects (auto parts, tires, lumber etc.). A general reference of composition was created, and our area of concern—large possibly harmful objects were accounted for. These items comprised the composition survey in 1968, 77', 78' and 79 to be 21.23%, 18.70%, 9.45%, and 10.44% consecutively out of all litter items. Once again, the amount and occurrence of undesirable objects (detrimental to the mowing operation) appears to be low enough to disregard as being of major concern.

Of the litter items studies in several cases, the litter items aluminum and paper products were found to be the greatest contributing factor. These items ranged from anywhere to ground level to approximately 4 inches in height. The items examined offer little resistance to the mowing blade.

Summary

Of the right-of-ways documented in the extensive literature search, it was determined that there is an infrequent, and rare occurrence of large, hazardous litter/trash objects along side the highways to cause rise of concern unless the mowing height was dropped below approximately 4 inches. This reduction in height which would encompass a larger portion of possibly harmful litters along side the highways. Should the optimum mowing height surpass a minimum of 3-4 inches, more concentration might then be placed on the amounts and quantity of litter items and or composition stratum.

References

- Albert, W. B. 1927. Studies on the growth of alfalfa and some perennial grasses. *J. Am. Soc. Agron.* 19:624-654.
- Barley, K. P. 1963. Influence of soil strength on growth of roots. *Soil Sci.* 95:175-180.
- Beard, J. B. 1973. *Turfgrass Science and Culture*. Prentice-Hall. Englewood Cliffs, NJ.
- Bell, C. C., and I. M. Ritchie. 1989. The effect of frequency and height of defoliation on the production and persistence of “ Grasslands Matua “ prairie grass. *Grass and Forage Science* 44: 245-248.
- Binnie, R. C., B. Chestnutt, and J. C. Murddoch. 1980. *Grass and Forage Science: J. British Grassl. Soc.* 35: 267-273.
- Biswell, H. H. and J. E. Weaver. 1933. Effect of frequent clipping on the development of roots and tops of grasses in prairie sod. *Ecology.* 14:368-390.
- Black, A. L., F. H. Siddoway, and R. W. Saulmon. (1971). “Wheatgrass Barriers Stop Soil Blowing, Trap Water.” *Montana Farmer-Stockman*, Vol 58. Pages 6 - 10.
- Borrelli, J., J. M. Gregory, and W. Abteu. (1989). “Wind Barriers: A Reevaluation of Height, Spacing, and Porosity.” *Transactions of the American Society of Agricultural Engineers*, Vol. 32, No. 6, pp. 2023 - 2027.
- Branson, F. A. 1956. Quantitative effects of clipping treatments of five range grasses. *J. Range Mgmt.* 9:86-88.
- Branson, F. A. 1953. Two factors affecting resistance of grasses to grazing. *J. Range Manage.* 6:165-171.
- Briske D. D. and J. H. Richards. 1995. Plant responses to defoliation: A physiologic, morphologic and demographic evaluation, p 635-710. *In*, Bedunah, D. J., and R. E. Sosebee E. (eds.). *Physiological ecology and developmental morphology*. (1st ed.). Soc. for Range Management, Denver, Colo.
- Briske, D. D., and J. H. Richards. 1994. Physiological responses of the individual plants to grazing, p. 147-176. *In*, M. Vavra, W. A. Laycock, and R. D. Pieper (eds.). *Ecological implications of livestock herbivory in the west*. (1st ed.). Soc. Range Management, Denver, Colo.
- Brown, F. Lee, and Lebeck, A. O. (1976). *Cars, Cans, and Dumps—Solutions for Rural Residuals*. Resources for the Future, The John Hopkins University Press, Baltimore,

Maryland.

Bruneau, A. H., J. E. Watkins and R. L. Brandenburg. 1992. Integrated pest management. p. 501-534. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) Turfgrass. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.

Bugge, W. A. and W. B. Snow. 1959. The complete highway. p. 3-32. *In* Snow, W. B. (Ed.) The Highway and the Landscape. Rutgers University Press, New Brunswick, NJ.

Bukey, F. S. and J. E. Weaver. 1939. Effects of frequent clipping on the underground food reserves of certain prairie grasses. *Ecology*. 20:246-252.

Burton, G. W., J. E. Jackson and R. H. Hart. 1963. Effects of cutting frequency and nitrogen on yield, in vitro digestibility, and protein fiber, and carotene content of coastal bermudagrass. *Agron. J.* 55:500-502.

Burzlaff, D. F., J. Stubbedieck, and J. Launchbaugh (no date). The growth and development of blue grama. Range and Wildlife Information Series. No. 1. Texas Tech University, Lubbock, Texas.

Busey, P. and J. H. Parker. 1992. Energy conservation and efficient turfgrass maintenance. p. 473-500. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) Turfgrass. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.

Busso, C. A., J. H. Richards, and N. J. Chatterton. 1989. Nonstructural carbohydrates and spring regrowth of two cool-season grasses: Interaction of drought and clipping. *J. Range Manage.* 43: 336-343.

Cable, D. R. 1982. Partial defoliation stimulates growth of Arizona cottontop. *J. Range Manage.* 35: 591-594.

Carter, J. F. and A. G. Law. 1948. The effect of clipping upon the vegetative development of some perennial grasses. *J. Am. Soc. Agron.* 40:1084-1091.

Clapp, J. G. Jr., D. S. Chamblee and H. D. Gross. 1964. Interrelationships between defoliation systems, morphological characteristics, and growth of Coastal bermudagrass. North Carolina Experiment Station and Agricultural Research Service, USDA. *Agron. Abs.*

Clark, P. E., W. C. Krueger, L. D. Bryant, and D. R. Thomas. 1998. Spring defoliation effects on bluebunch wheatgrass. II. Basal area. *J. Range Manage.* 51: 526-530.

Cogburn, B. 1995. TxDOT Mowing Guidelines. Native Plant Society of Texas Vol.

XIII, No.3.

Coyne, P. I., And C. W. Cook. 1970. Carbohydrate reserve cycles in eight desert range species. *J. Range Manage.* 23: 438-444.

Coyne, P. I., M. J. Trlica, and C. E. Owensby. 1995. Carbon and nitrogen dynamics in range plants, p. 59-167. *In*, D. J. Bedunah, and R. E. Sosebee (eds.) *Physiological ecology and developmental morphology.* (1st.ed). Soc. Range Management. Denver, Colo.

Crider, F. J. 1955. Root growth stoppage resulting from defoliation of grass. U.S. Tech. Bull. No. 1102. pp. 1-23.

Crone, F. W. 1959. The art of fitting the highway to the landscape. p. 78-109. *In* Snow, W. B. (Ed.) *The Highway and the Landscape.* Rutgers University Press, New Brunswick, NJ.

Crovetto, C. 1996. *Stubble over the Soil.* Am. Soc. Agron. Madison, WI.

Dahl, B. E. 1995. Developmental morphology and management implications, p. 22-58. *In*, D. J. Bedunah, and R. E. Sosebee (eds.). *Physiological ecology and developmental morphology.* (1st ed.). Soc. Range Mangement. Denver, Colo.

Dahl, B. E., and D. N. Hyder. 1977. Developmental morphology and management implications, p. 257-290. *In*, R. E. Sosebee (ed.) *Rangeland plant physiology.* Range Sci. Series No. 4. Soc. For Range Mange. Denver, Colo.

Darryl, G. S., A. McLean, B. Brooke, and J. Hall. 1980. Influence of simulated grazing (clipping) on pinegrass growth. *J. Range Manage.* 33: 286-291.

Davis, R. R. 1967. Population changes in Kentucky bluegrass-red fescue mixtures. Ohio Agr. Res. and Dev. Center. Agron. Abs.

DiPaola, J. M. and J. B. Beard. 1992. Physiological effects of temperature stress. p. 231-268. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) *Turfgrass.* Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.

Dodd, J. D. and H. H. Hopkins. 1958. Yield and carbohydrate content of blue grama grass as affected by clipping. *Trans. Kansas Academy Sci.* 61: 280-287.

Donart, G. B., and C. W. Cook. 1970. Carbohydrate reserves content of montain range plants following defoliation and regrowth. *J. Range Manage.* 23: 15-19.

Duble, R. L. 1996. *Turfgrasses: Their Management and Use In The Southern Zone* Texas A&M University Press. College Station, TX.

El Hassan, B., and W. C. Krueger. 1980. Impact of intensity and season of grazing on carbohydrate reserves of perennial ryegrass. *J. Range Manage.* 33: 200-204.

Everson, A. C. 1966. Effects of frequent clipping at different stubble heights on western wheatgrass (*Agropyron smithii* Rybd.). *Agron. J.* 58:33-35.

Finkner, A. L. (1969). *National Study of the Composition of Roadside Litter*. Report prepared for To Keep America Beautiful, Inc., Highway Research Board of the Division of Engineering, National Research Council-National Academy of Sciences – National Academy of Engineering. New York.

Fluck, R. C. and P. Busey. 1998. Energy for mowing turfgrass. *Trans. ASAE* 31: 1304-1308.

Follett, R. H., L. S. Murphy and R. L. Donahue. 1981. *Fertilizers and Soil Amendments*. Prentice-Hall, Englewood Cliffs, NJ.

Forwood J. R., and M. M. Magai. 1992. Clipping frequency and intensity effects on big bluestem yield, quality, and persistence. *J. Range Manage.* 45: 554-559.

Gerard, C. J., H. C. Mehta and E. Hinojosa. 1972. Root growth in a clay soil. *Soil Sci.* 14:37-49.

Gillen, R. L., and R. W. McNew. 1987. Seasonal growth rates of tallgrass prairie after clipping. *J. Range Manage.* 40: 342-345.

Goss, R. L. and A. G. Law. 1967. Performance of bluegrass varieties at 2 cutting heights and 2 nitrogen levels. *Agron. J.* 59:516-518.

Graber, L. F. and H. W. Ream. 1931. Growth of bluegrass with various defoliations and abundant nitrogen supply. *J. Am. Soc. Agron.* 23:938-946.

Greb, B. W. (1980). *Snowfall and Its Potential Management in the Semiarid Central Great Plains*. ARM-W-18, U.S. Department of Agriculture, Science and Education Administration Reviews and Manuals. 46 pages.

Greeley, Ronald, and Iversen, James D. (1985). *Wind as a Geological Process on Earth, Mars, Venus, and Titan*. Cambridge University Press, Cambridge, England. 333 pages.

Haferkamp, M. R. 1982. Defoliation impacts on quality and quantity of forage harvested from sacaton. *J. Range Manage.* 35: 26-31.

Hagen, L. J., and Skidmore, E. L. (1971). "Turbulent Velocity Fluctuations and Vertical Flow as Affected by Windbreak Porosity." *Transactions of the American Society of Agricultural Engineers*, Vol. 14, No. 3, pp. 634 - 637.

Harivandi, M. A., J. D. Butler and L. Wu. 1992. Salinity and turfgrass culture. p. 207-230. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) Turfgrass. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.

Harrison, C. M. 1931. Effect of cutting and fertilizer application on grass development. *Plant Phys.* 6:669-684.

Harrison, C. M. 1931. Effect of cutting and fertilizer applications on grass development. *Plant Phys.* 6:669-682.

Harrison, C. M. 1934. Responses of Kentucky bluegrass to variations in temperature, light, cutting, and fertilizing. *Plant Phys.* 9:83-106.

Harrison, C. M. and C. W. Hodgson. 1939. Response of certain perennial grasses to cutting treatments. *J. Am. Soc. Agron.* 31:418-430.

Hart, R. H. 1974. Crop selection and management. *In* Factors Involved in Land Application of Agricultural and Municipal Wastes. USDA-ARS. Beltsville, Md.

Hayes, M. J., and E. D. Williams. 1995. Effects of timing and intensity of defoliation on growth and development of white clover seedlings. *Grass and Forage Science.* 50: 414-423.

Hendrickson A. H. and F. J. Veihmeyer. 1931. Influence of dry soil on root extension. *Plant. Phys.* 6:567-576.

Heppner, M. B. 1961. No summer dormancy. *The Farm Quart.* 16: 1-7.

Herford, L.R. 1951. The effect of different intensities and frequencies of clipping on forage yield of *Andropogon scoparius* Michx. and *Paspalum plicatulum* Michx. M.S. Thesis. Texas A&M University. College Station.

Hill, M., and R. W. Watson. 1989. The effect of differences in intensity and frequency of defoliation on the growth of phalaris and cocksfoot. *Aus. J. Agr. Res.* 40: 333-343.

Hudson, N. 1995. *Soil Conservation*, 3rd ed. Iowa State University Press. Ames, IA.

Hughes, H. D. 1937. Response of *Poa pratensis* L. to different harvest treatments, measured by weight and composition of forage and roots. Report of the 4th Int. Grassland Congr. Aberystwyth, Wales. pp.447-452.

Hull, R. J. 1992. Energy relations and carbohydrate partitioning in turfgrasses. p. 175-206. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) Turfgrass.

Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.

Hunt, Roy E., and Deschamps, Richard J. (1995). "Stability of Slopes." In *The Civil Engineering Handbook* edited by W. F. Chen. CRC Press, New York. 2609 pages.

Hyder, D. N. 1972. Defoliation in relation to vegetative growth, p. 304-317. In, V. B. Younger, and C. M. McKell (eds.). *Biology and utilization of grasses*. Academic press, New York.

Hyder, D. N. 1974. Morphogenesis and management of perennial grasses in the U.S., p. 89-98. In, *Plant morphogenesis as the basis for scientific management of range resources*. U. S./Australia rangelands Workshop Proc. Berkeley, Calif. U.S. Dep. Agr. Misc. Publ. 1271.

Ingraham, J. C. 1959. Politics and road building. p. 169-181. In Snow, W. B. (Ed.) *The Highway and the Landscape*. Rutgers University Press, New Brunswick, NJ.

Jameson, D. A., and D. L. Huss. 1958. The effect of clipping leaves and stems on number of tillers, herbage weights, root weights, and food reserves of little bluestem. *J. Range Manage.* 12: 122-126.

Johnson, W. A. 1959. Preserving the scenic qualities of the roadside. p. 110-130. In Snow, W. B. (Ed.) *The Highway and the Landscape*. Rutgers University Press, New Brunswick, NJ.

Juska, F. V. and A. A. Hanson. 1961. Effects of interval and height of mowing on growth of Merion and common Kentucky Bluegrass (*Poa Pratensis* L). *Agron. J.*

Kalmbacher, R. S., F.G. Martin, and J. M. S. Andrade. 1981. Yield and quality of creeping bluestem as affected by time of cutting. *J. Range Manage.* 34: 471-474.

Keep Australia Beautiful Council. (1971). *A Study of Roadside Litter*. A report by the Institute of Applied Economic and Social Research, University of Melbourne, for the Keep Australia Beautiful Council, Melbourne, Victoria. 8 pages.

Kneebone, W. R. 1966. Root growth of bermudagrass varieties at three mowing heights in 1966. *Arizona Ag. Expt. Sta. Report* 240:15-17.

Kneebone, W. R., D. M. Kopec and C. F. Mancino. 1992. Water requirements and irrigation. p. 441-472. In Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) *Turfgrass*. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.

Kowalenko, B. L., and J. T. Romo. 1998. Regrowth and rest requirements of northern wheatgrass following defoliation. *J. Range Manage.* 51: 73-78.

- Krenisky, E. C., M. J. Carroll, R. L. Hill and J. M. Krouse. 1998. Runoff and sediment losses from natural and man-made erosion control materials. *Crop Sci.* 38:1042- 1046.
- Lacey, J. R., K. M. Olson-Rutz, M. Haferkamp, and G. A. Kennett. 1994. Effects of defoliation and competition on total non-structural carbohydrates of spotted knapweed. *J. Range Manage.* 47: 481-484.
- Larson, W. E., F. J. Pierce and R. H. Dowdy. 1983. The threat of soil erosion to long-term crop productivity. *Science* 219:458-465.
- Lattanzi, A. R., L. D. Meyer and M. F. Baumgardner. 1979. Influence of mulch rate and slope steepness in interrill erosion. *Soil Sci. Soc. Am. Proc.* 38:946-950.
- Law, A. G. 1966. Performance of bluegrass varieties clipped at two heights. *Weeds, Trees and Turf.* 5:26-28.
- Lowe, K. F., and T. M. Bowdler. 1988. Effects of height and frequency of defoliation on the productivity of irrigated oats and perennial ryegrass, grown alone or with barrel medic. *Aus. J. Exp. Agr.* 28: 57-67.
- Madison, J. H. 1971. *Practical Turfgrass Management.* Van Nostrand Reinhold Co. New York, NY. p. 96-124.
- Madison, J. H. 1932. Effects of mowing, irrigation, and nitrogen treatment of *Agrostis palustris* Huds., "Seaside" and *Arostis tenuis* Sibth., "Highland" on population, yield, rooting, and cover. *Agron. J.* 54:407-412.
- McCarty, E. C. and R. Price. 1942. Growth and carbohydrate content of important mountain forage plants in central Utah affected by clipping and grazing. USDA Tech. Bull. No. 818. pp. 1-51.
- McCully, W. G. and W. J. Bowmer. 1967. Promoting establishment of vegetation for erosion control. TxDOT. Res. Project 2-18-63-67. Res. Rept. 67-6.
- McCully, W. G. and W. J. Bowmer. 1969. Erosion control on roadsides in Texas. TxDOT. Res. Project 2-18-63-67. Res. Rept. 67-8. (Final)
- McCully, W. G. and W. J. Bowmer. 1971. Evaluation of soil sterilant herbicides for roadsides. TxDOT. Res. Project 2-18-69-142. Res. Rept. 142-2.
- Meier, G. and T. Weaver. (Eds.) 1997. Desirables and weeds for roadside management—a Northern rocky Mountain catalogue. US Dept. Trans. Res. Development, and Technology Transfer Program. MS-7069 TTS 98-8 #30.

Menke, J. W., and M. J. Trlica. 1981. Carbohydrate reserve, phenology, and growth cycles of nine Colorado range species. *J. Range Manage.* 34: 269-277.

Meredith H. L. and W. H. Patrick, Jr. 1961. Effects of soil compaction on subsoil root penetration and physical properties of three soils in Louisiana. *Agron. J.* 66:341- 344.

Miller, R. F., M. R. Haferkamp, and R. F. Angel. 1990. Clipping date effects on soil water and regrowth in crested wheatgrass. *J. Range Manage.* 43: 253-357.

Mitchell, K. J. 1955. Growth of pasture species. II. Perennial ryegrass (*Lolium perenn*), cocksfoot (*Dactylis glomerate*), and paspalum (*Paspalum dilatatum*). *New Zealand J. Sci. Tech. Sec. A.* 37:8-26.

Mortimer, G. B. and H. L. Ahlgren. 1936. Influence of fertilization, irrigation, and stage and height of cutting on yield and composition of Kentucky bluegrass. *J. Am. Soc. Agron.* 28:515-533.

Moisy, E. B., and McPherson, F. B. (1966). "Effect of Porosity on Performance of Windbreaks." *Transactions of the American Society of Agricultural Engineers.* Vol. 9, No. 1, pp 74 - 76.

Mullahey, J. J., S. S. Waller, and L. E. Moser. 1990. Defoliation effects on production and morphological development of little bluestem. *J. Range Manage.* 43: 497-501.

Mullahey, J. J., S. S. Waller, and L. E. Moser. 1991. Defoliation effects on yield and bud and tiller numbers of two sandhills grasses. *J. Range Manage.* 44: 241-245.

Murphy, J. S., and D. D. Briske. 1992. Regulation of tillering by apical dominance: Chronology, interpretive value, and current perspectives. *J. Range Manage.* 45: 419-429.

O'Toole, M. L. (1979). *Michigan Roadside Litter Composition Study.* Michigan Department of Transportation, Lansing, Michigan. 28 pages.

Pennsylvania State University. 1968. Dept. Of Agricultural Education. Turfgrass Maintenance and Establishment: A Student Handbook. University Park, PA.

Phillips, R. E. and D. Kirkham. 1962. Mechanical impedance and corn seedling root growth. *Soil Sci. Soc. Am. Proc.* 26: 319-322.

Portas, C. A. M. and H. M. Taylor. 1976. Growth and survival of young plant roots in dry soil. *Soil Sci.* 121:170-175.

Problems Related to Windbreaks. SCS-TP-112, United States Department of Agriculture, Soil Conservation Service, Washington, D.C., 25 pages.

Pugh, H. L. D., and Price, W. I. J. (1954). "Snow Drifting and the Use of Snow Fences." *Polar Record*, Vol. 7, No. 47, pp. 4 - 23.

Raine, J. K., and Stevenson, D. C. (1977). "Wind Protection by Model Fences in Simulated Atmospheric Boundary Layer." *Journal of Industrial Aerodynamics*, Vol. 2, page 159 - 180.

Razmjoo, K., T. Imada, J. Suguira and S. Kaneko. 1996. Effect of nitrogen rates and mowing heights on color, density, uniformity, and chemical composition of creeping bentgrass cultivars in winter. *J. Plant Nut.* 19:1499-1509.

Rechenthin, C. A. 1956. Elementary morphology of grass growth and how it effects utilization. *J. Range Manage.* 9:167-170.

Richards, J. H., R. J. Mueller, and J. J. Mott. 1988. Tillering in tussock grasses in relation to defoliation – apical bud removal. *Ann. Bot.* 62:173-179.

Robertson, J. H. 1933. Effect of frequent clipping on the development of certain grass seedlings. *Plant Phys.* 8:425-447.

Salaiz, T. A., Horst, G. L. and R. C. Shearman. 1995. Mowing height and vertical mowing frequency effects on putting green quality. *Crop Sci.* 35: 422-442.

Schroeder, C. B. 1996. *Turf Management Handbook*. Interstate Press. Danville, IL.

Schuurman, J. J. 1965. Influence of soil density on root development and growth of oats *Plant Soil* 22:352-374.

Schwab, G. O., D. D. Fangmeier and W. J. Elliot. 1996. *Soil and Water Management Systems*, 4th ed. Wiley. New York, NY.

Skidmore, E. L. and F. H. Siddoway. 1978. Crop residue requirements to control wind erosion. *In* W. R. Oschwalk. (Ed.) *Crop Residue Management Systems*. ASA Special Publ. # 31. Am. Soc. Agron., Crop Sci. Soc. Am. and Soil Sci. Soc. Am. Madison, WI.

Skidmore, E. L., and Hagen, L. J. (1977). "Reducing Wind Erosion with Barriers." *Transactions of the American Society of Agricultural Engineers*, Vol. 20, No. 3. Pages 911 - 915.

Snow, W. B. 1959. *The Highway and the Landscape*. Rutgers University Press, New Brunswick, NJ.

Sprague, H. B. 1933. Root development of perennial grasses and its relation to soil conditions. *Soil Sci.* 36:189-209.

Syrek, Daniel B. (1986). *TEXAS LITTER: 1985 – 1986—The Change is Visible Litter Rates at 100 State Highway Locations*. Report prepared for The Texas Department of Highways and Public Transportation. The Institute for Applied Research, Sacramento, California. 39 pages.

Syrek, Daniel B. (1989). *California Litter: One Year After—The Impact of the Beverage Container Recycling and Litter in California*. Report prepared for the State of California, Department of Conservation, Division of Recycling. The Institute for Applied Research, Sacramento, California. 52 pages.

Syrek, Daniel B. (1990). *LOUISIANA LITTER: 1990—A Survey of Visible Litter*. Report prepared for The Louisiana Department of Environmental Quality. The Institute for Applied Research, Sacramento, California. 44 pages.

Tabler, Ronald D. (1980). "Geometry and Density of Drifts Formed by Snow Fences." *Journal of Glaciology*, Vol. 26, No. 94, pp. 405 - 419.

Taylor, H. M. and E. Burnett. 1964. Influence of soil strength on the root-growth habits of plants. *Soil Sci.* 98:174-180.

Texas Department of Transportation. 1993. *Roadside Vegetation Management Manual*. A volume of the *Infrastructure Maintenance Manual*. Manual Notice 93-1.

Texas Department of Transportation. 1996. *Roadside Vegetation Management Manual*. Manual Notice 96-1.

Texas Department of Transportation. 1999. *Giving Nature a Hand*. booklet.

Texas Transportation Institute. 1996. Weed control geotextile tried in Texas. *Better Roads*. 66(10):23-24.

Thompson, D. B. and R. E Zartman. 1997. Soil erosion along Interstate highway 20 near Longview, TX. *Proc. Int. Erosion Control Assn.*

Tisdale, S. L. and W. L. Nelson. 1975. *Fertility and Fertilizers*. 3rd Ed. Macmillan Publishing Co., New York, NY.

Transportation Research Board. 1974. Highway deicing comparing alt and calcium magnesium acetate. National Research Council. Special Report 235.

Trlica, M. J. 1977. Distribution and utilization of carbohydrate reserves in range plants, p. 73-96. *In*, R. E. Sosebee (ed.). *Rangeland plant physiology*. Range Sci. Series No. 4. Soc. Range Management. Denver, Colo.

Trlica, M. J., and C. W. Cook. 1971. Defoliation effects on carbohydrate reserves of desert species. *J. Range Manage.* 24: 418-425.

- Troeh, F. R. and L. M. Thompson. 1993. *Soils and Soil Fertility*. 5th Ed. Oxford Univ. Press. New York, NY.
- Troeh, F. R., J. A. Hobbs and R. L. Donahue. 1980. *Soil and Water Conservation for Productivity and Environmental Protection*. Prentice-Hall, Englewood Cliffs, NJ.
- Turgeon, A. J. 1999. *Turfgrass Management*. 5th Ed. Penn State University. Prentice Hall, Upper Saddle River, NJ.
- Turner, T. R. and N. W. Hummel, Jr. 1992. Nutritional requirements and fertilization. p. 385-440. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) *Turfgrass*. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.
- Undersander, D. J., and C. H. Naylor. 1987. Influence of clipping frequency on herbage yield and nutrient content of tall wheatgrass. *J. Range Manage.* 40: 31-35.
- Vesilind, P. A. (1976). *Measurement of Roadside Litter*. Duke Environmental Center, Duke University. Durham, N.C. 15 pages.
- Vogel, W. G., and A. J. Bjustad. 1968. Effects of clipping on yield and tillering of little bluestem, big bluestem, and Indiangrass. *J. Range Manage.* 21: 136-140.
- Waddington, D. V., R. N. Carrow and R. C. Shearman. 1992. (Eds.) *Turfgrass*. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.
- Watschke, T. L. and R. E. Schmidt. 1992. Ecological aspects of turf communities. p. 129-174. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) *Turfgrass*. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.
- Watschke, T. L., M. G. Prinster and J. M. Breuninger. 1992. Plant growth regulators and turfgrass management. p. 557-588. *In* Waddington, D. V., R. N. Carrow and R. C. Shearman. (Eds.) *Turfgrass*. Agronomy monograph #32. Am. Soc. Agron, Crop Sci. Soc. Am., and Soil Sci. Soc. Am. Madison, WI.
- White, E. M. and J. K. Lewis. 1969. Ecological effect of a clay soils structure on some native grass roots. *J. Range Mgmt.* 22:401-404.
- White, R. P. 1959. The functional uses of plants on the complete highway. p. 182-192. *In* Snow, W. B. (Ed.) *The Highway and the Landscape*. Rutgers University Press, New Brunswick, NJ.
- Willms, W. D. 1991. Cutting frequency and cutting height effects on rough fescue and parry oat grass yield. *J. Range Manage.* 44: 82-86.

Woodruff, N. P., and Zingg, A. W. (1952). Wind-Tunnel Studies of Fundamental
Zhang, J., and J. T. Romo. 1994. Defoliation of a northern wheatgrass community:
Above- and below ground phytomass productivity. *J. Range Management*.